

Associations between eating speed, diet quality, adiposity, and cardiometabolic risk factors

Tany E. Garcidueñas-Fimbres, MD, MSc^{1,2,3}, Indira Paz-Graniel, PhD^{1,2,3}, Carlos Gómez-Martínez, MSc^{1,2,3}, Jose Manuel Jurado-Castro, PhD^{1,4}, Rosaura Leis, MD, PhD^{1,5,6,7}, Joaquin Escribano, MD, PhD^{3,8}, Luis A. Moreno, MD, PhD^{1,9,10}, Santiago Navas-Carretero, PhD^{1,11,12}, Olga Portoles, PhD^{1,13}, Karla A. Pérez-Vega, MSc^{1,14}, Mercedes Gil-Campos, MD, PhD^{1,4}, Alicia López-Rubio, MSc^{5,6,7}, Cristina Rey-Reñones, PhD^{15,16,17}, Pilar De Miguel-Etayo, PhD^{1,9,10}, J. Alfredo Martínez, MD, PhD¹¹, Katherine Flores-Rojas, PhD⁴, Rocío Vázquez-Cobela, PhD^{1,5,6,7}, Verónica Luque, PhD^{3,8}, Maria Luisa Miguel-Berges, PhD^{1,9,10}, Belén Pastor-Villaescusa, PhD⁴, Francisco Jesus Llorente-Cantarero, PhD^{1,18}, Jordi Salas-Salvadó, MD, PhD^{1,2,3,*}, and Nancy Babio, PhD^{1,2,3,*}, on behalf of the Childhood Obesity Risk Assessment Longitudinal Study (CORALS) study investigators

Objective To assess the associations between eating speed, adiposity, cardiometabolic risk factors, and diet quality in a cohort of Spanish pre-school-children.

Study design A cross-sectional study in 1371 preschool age children (49% girls; mean age, 4.8 ± 1.0 years) from the Childhood Obesity Risk Assessment Longitudinal Study (CORALS) cohort was conducted. After exclusions, 956 participants were included in the analyses. The eating speed was estimated by summing the total minutes used in each of the 3 main meals and then categorized into slow, moderate, or fast. Multiple linear and logistic regression models were fitted to assess the β -coefficient, or OR and 95% CI, between eating speed and body mass index, waist circumference, fat mass index (FMI), blood pressure, fasting plasma glucose, and lipid profile.

Results Compared with participants in the slow-eating category, those in the fast-eating category had a higher prevalence risk of overweight/obesity (OR, 2.9; 95% CI, 1.8-4.4; $P < .01$); larger waist circumference (β , 2.6 cm; 95% CI, 1.5-3.8 cm); and greater FMI (β , 0.3 kg/m²; 95% CI, 0.1-0.5 kg/m²), systolic blood pressure (β , 2.8 mmHg; 95% CI, 0.6-4.9 mmHg), and fasting plasma glucose levels (β , 2.7 mg/dL, 95% CI, 1.2-4.2 mg/dL) but lower adherence to the Mediterranean diet (β , -0.5 points; 95% CI, -0.9 to -0.1 points).

Conclusions Eating fast is associated with higher adiposity, certain cardiometabolic risk factors, and lower adherence to a Mediterranean diet. Further long-term and interventional studies are warranted to confirm these associations. (*J Pediatr* 2022; ■:1-9).

The vast majority of strategies to prevent or treat overweight and obesity in children are based on increasing physical activity, decreasing sedentary behaviors, and promoting adherence to a healthy diet.¹ However, some

From the ¹Consorcio CIBER, Fisiopatología de la Obesidad y Nutrición, Instituto de Salud Carlos III, Madrid, Spain; ²Universitat Rovira i Virgili, Departament de Bioquímica i Biotecnologia, Unitat de Nutrició Humana, Reus, Spain; ³Institut d'Investigació Sanitària Pere Virgili, Reus, Spain; ⁴Reina Sofia University Hospital, Maimónides Biomedical Research Institute of Córdoba, University of Córdoba, Córdoba, Spain; ⁵Unit of Pediatric Gastroenterology, Hepatology and Nutrition, Pediatric Service, Complejo Hospitalario Universitario de Santiago de Compostela, Santiago de Compostela, Spain; ⁶Pediatric Nutrition Research Group, Health Research Institute of Santiago de Compostela, Santiago de Compostela, Spain; ⁷Unit of Investigation in Human Nutrition, Growth and Development of Galicia, University of Santiago de Compostela, Santiago de Compostela, Spain; ⁸Pediatrics, Nutrition, and Development Research Unit, Universitat Rovira i Virgili, Reus, Spain; ⁹Growth, Exercise, Nutrition and Development Research Group, Instituto Agroalimentario de Aragón, Instituto de Investigación Sanitaria Aragón, University of Zaragoza, Zaragoza, Spain; ¹⁰Department of Psychiatry and Nursing, University of Zaragoza, Zaragoza, Spain; ¹¹Center for Nutrition Research, University of Navarra, Pamplona, Spain; ¹²IdiSNA Institute for Health Research, Pamplona, Spain; ¹³Department of Preventive Medicine and Public Health, University of Valencia, Valencia, Spain; ¹⁴Cardiovascular Risk and Nutrition Research Group, Hospital del Mar Medical Research Institute, Barcelona, Spain; ¹⁵Departament de Ciències Mèdiques Bàsiques, Facultat de Medicina, Universitat Rovira i Virgili, Reus, Spain; ¹⁶Unitat de Suport a la Recerca Camp de Tarragona, Fundació Institut Universitari per a la Recerca a l'Atenció Primària de Salut Jordi Gol i Gurina, Reus, Spain; ¹⁷Centre d'Atenció Primària St Pere, Direcció d'Atenció Primària Camp de Tarragona, Institut Català de la Salut, Reus, Spain; and ¹⁸Department of Specific Didactics, Faculty of Education, University of Córdoba, Córdoba, Spain

*Co Senior authors and contributed equally.

A list of additional members of the CORALS study investigators is available at www.jpeds.com (Appendix).

Supported by the Consejo Nacional de Ciencia y Tecnología (769789, to T.G.-F.); the Spanish Ministry of Education, Culture, and Sports (FPU 17/01925, to I.P.-G.); and the Universitat Rovira i Virgili (2020PMF-PIPF-37, to C.G.-M.). V.L. received a Serra Hunter Fellowship from Generalitat de Catalunya. J.S.-S. is partially supported by the Catalan Institute for Research and Advanced Studies (ICREA) under the ICREA Academia program. Funds for the establishment of the CORALS cohort in the first year of the study (2019) were provided by an agreement between the Danone Institute and the Spanish Biomedical Research Center on Physiopathology of Obesity and Nutrition. The authors declare no conflicts of interest.

0022-3476/© 2022 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). <https://doi.org/10.1016/j.jpeds.2022.08.024>

BMI	Body mass index
CEBQ	Child Eating Behaviour Questionnaire
CESNID	Centre d'Ensenyament Superior de Nutrició i Dietètica
CORALS	Childhood Obesity Risk Assessment Longitudinal Study
FMI	Fat mass index
HDL-c	High-density lipoprotein cholesterol
LDL-c	Low-density lipoprotein cholesterol

recent studies have reported that greater adiposity and other cardiometabolic risk factors also can be influenced by certain behaviors, such as eating speed and eating frequency, among others, in adults, adolescents, and children.² Eating quickly has been related to higher dietary energy intake, body mass index (BMI),² obesity prevalence,^{3,4} and certain metabolic disorders.^{2,5} In contrast, chewing slowly and increasing the number of chewing cycles during a meal have been negatively associated with adiposity.⁶ It has been proposed that a slower eating speed might enhance the development of a satiety signal, which could limit the total food intake.⁷

Even though a universal definition for “speed of eating” has not been settled, a few authors define it as the time that a person takes to eat one meal.⁸ Other authors refer to “eating rate” (as the estimation in grams per minute of food consumed)⁹ or “energy intake rate” (energy intake by unit of time; kcalories per minute),¹⁰ which may be considered more objective measures. Eating rate has been associated with food textures,¹¹ tastes,¹⁰ and an increased overall energy.¹² Some authors have suggested that a fast eating rate is associated with high-fat content meals,¹⁰ which usually have a softer texture and intense taste.¹³ In addition, the consumption of soft-textured foods may be associated with a lower number of chewing cycles.¹⁴ Meanwhile, a low energy intake rate has been associated with solid textures and low water content,¹⁰ whereas a greater energy intake rate has been related to fat, sweet, and sour taste intensity,¹⁰ as well as to increased consumption of ultra-processed foods that may contribute to poor dietary quality.¹⁵ A high energy intake rate also has been associated with a faster eating rate.¹⁶

To our knowledge, associations between eating speed and diet quality have not been explored in children, and their associations with cardiometabolic risk factors have been rarely assessed. The present cross-sectional study aimed to assess the associations between eating speed, diet quality, adiposity, and cardiometabolic risk factors in a Spanish cohort of children aged 3-6 years.

Methods

This cross-sectional study is based on the baseline data of the Childhood Obesity Risk Assessment Longitudinal Study (CORALS), a prospective ongoing multicenter study conducted in preschool children that has a 10-year expected follow-up (<https://corals.es/>). The main aim of the CORALS project is to identify potential risk factors for childhood obesity. Briefly, eligible participants are children aged 3-6 years attending the selected schools across 7 Spanish cities (Barcelona, Córdoba, Pamplona, Reus, Santiago de Compostela, Valencia, and Zaragoza) whose tutors had agreed to participate. To be enrolled in the study, parents or caregivers had to sign a consent form, attend the inclusion visit, and complete several questionnaires (eg, leisure time physical activity, 3-day food record, social-demographic data). The exclusion criteria include belonging to a family with difficulty participating, comprehension or language difficulties, and unstable residence.

The Ethics Committee of each recruitment center approved the study protocol (reference nos. 051/2019, 4155/2019, 2019/18, 9/19, 09/2019, 19/27, and 2019/131), which was conducted following the standards of the Declaration of Helsinki.

Children aged 3-6 years recruited between March 2019 and June 2021 were selected (n = 1371). Participants who attended the baseline visit and completed the provided questionnaires, as well as those who had available data in the duration of breakfast, lunch, and dinner, were included in the present analyses. Children with current diagnosis of chronic diseases, such as diabetes mellitus type 2, hypertension, and familiar hypercholesterolemia, were excluded from these analyses.

Eating speed was assessed through the question “How long does it usually take for your child to eat in each meal?” Speed was calculated by summing the total time (in minutes) reported by parents in the 3 main meals (breakfast, lunch, and dinner). Then categories for slow, moderate, and fast-eating speed were determined by tertiles. These categories were recategorized so that the highest tertile represents the fast-eating category. Additional analyses were performed using as exposure variable the “slowness in eating” scale from the validated Child Eating Behaviour Questionnaire (CEBQ),¹⁷ which has been recently validated in Spanish children of the same age.¹⁸ The possible responses included on the CEBQ “slowness in eating” scale are “my child finishes his/her meal quickly,” “my child eats slowly,” “my child takes more than 30 minutes to finish a meal,” and “my child eats more and more slowly during the course of a meal.” This scale was categorized by tertiles that were recategorized, with the third tertile as the fast-eating category.

Body weight (in kilograms) and body fat mass (in kilograms) were assessed with a precision scale (Tanita MC780SMA; Tanita Europe, B.V.). Body fat mass was assessed by octopolar multifrequency bioelectrical impedance. Height (in centimeters) was assessed using a portable stadiometer (seca 213, Escala 20-205 cm; SECA). These assessments were performed with the child in light clothing, without shoes, and according to standard procedures. BMI was calculated as weight (in kilograms) divided by height (in meters squared) and classified as underweight/normal weight or overweight/obesity using the cutoff points defined by Cole et al.¹⁹ Waist circumference was determined with a measuring tape (seca 201), midway between the lowest rib and the iliac crest. The fat mass index (FMI) was estimated as body fat mass (in kilograms)/squared height (in meters).²⁰

Blood pressure was determined in the nondominant arm and measured 3 times (with 5-minute intervals between measurements) using an automatic oscillometer (Omron M3 Intellisense HEM-75051-EV; IOMRON Healthcare Europe B.V.) with a child-adapted cuff. Blood pressure was recorded as the mean of the 3 measurements for both systolic and diastolic blood pressures.

Blood samples were collected after a minimum of 8 hours of fasting. Serum samples were used to determine levels of fasting plasma glucose, total cholesterol, high-density

lipoprotein cholesterol (HDL-c), low-density lipoprotein cholesterol (LDL-c), and triglycerides by standard procedures. Non-HDL-c was calculated with the equation suggested by Frost et al.²¹

Diet quality was assessed by an ad hoc questionnaire of 18 items adapted to children from a 14-item Mediterranean diet questionnaire validated in an elderly population.²² This questionnaire evaluates the consumption of typical foods in the Mediterranean diet (eg, olive oil, fruits, vegetables, cereals, nuts, seeds, cheese, yogurt, legumes, fish, seafood, poultry, rabbit), as well as additional foods that are not included in this dietary pattern (ie, butter, cream, carbonated/sugary beverages, junk food, dairy desserts, and precooked/ready-to-eat foods). The number of servings has been adapted for several items, and certain typical foods consumed by children (eg, gummy worms, sweets, petit suisse) are also included in this questionnaire (**Table I**; available at www.jpeds.com). Compliance with each of the 18 items was scored as 1 point; thus, the total score ranged from 0 to 18, with 0 representing no adherence and 18 representing the greatest adherence.

A set of self-administered questionnaires completed by parents or caregivers at home was used to collect information on early life factors, maternal data (eg, age, BMI, educational level, socio-professional status), and lifestyle patterns, among others. Weight at birth (kilograms), mother's weight gain during pregnancy (kilograms), and breastfeeding (months of age) were recorded.

A physical activity questionnaire based on the Outdoor Playtime Checklist and the Outdoor Playtime Recall Questions²³ was provided. An additional ad hoc questionnaire with 13 components based on leisure physical activity that includes sedentary behaviors was used to determine the active lifestyle score (**Table II**; available at www.jpeds.com).

Sleep pattern was explored with the questions "how long does your child sleep at night during weekdays and on weekend days or holidays?" and "how long does your child nap during weekdays and on weekend days or holidays?" The total sleep duration (hours per day) was calculated summing night sleep hours and nap time on weekdays and weekend days or holidays divided by 7 (ie, the total days of the week).

Trained dietitians assessed the dietary intake using a semi-quantitative 125-item food frequency questionnaire adapted from a food frequency questionnaire validated in adults.²⁴ The 9 possible answers ranged from "never" to "more than 6 times per day" and were transformed into grams or milliliters per day using the standard portion size of each item. The CESNID (Centre d'Enseignement Supérieur de Nutrición i Dietètica) database was used to calculate total energy and nutrient intake.²⁵ Total energy intake was estimated according to Goldberg cutoffs adapted to children²⁶; participants with missing data or implausible reported energy intake were excluded from the analyses.

The present analyses were conducted using the CORALS database updated to December 2021. To compare general characteristics and dietary variables among categories of eating speed, one-factor ANOVA, the Kruskal–Wallis test, or χ^2 test

was used. The data were presented as mean \pm SD or median (IQR) for quantitative variables and as percentage (number) for qualitative variables. The Kolmogorov–Smirnov test was used to assess the normal distribution of variables. Missing data <10% on covariates was imputed to the mean.²⁷ Multiple linear regression models were fitted to assess associations between eating speed (exposure) and different outcomes—waist circumference, FMI, systolic blood pressure, diastolic blood pressure, fasting plasma glucose, total cholesterol, HDL-c, LDL-c, non-HDL-c, triglycerides, and Mediterranean diet score—which were reported as β coefficients and 95% CIs. Logistic regression models were fitted to explore the association between eating speed and weight status (outcome), expressed as OR and 95% CIs. All models were adjusted by the following confounders: center, sex, age, maternal BMI, and mother's educational level, total energy intake, breastfeeding, total minutes of physical activity a week, and total hours of sleep per day. In our analyses, the slower category of eating speed served as the reference. Interaction analyses were performed by sex, maternal factors (BMI and educational level), weight at birth, and sedentary behaviors.

Sensitivity analyses were performed for FMI excluding those participants aged 3–4 years owing to a lack of validation of bioimpedance equations (Tanita MC780SMA) in children aged <5 years. Correlation analyses were performed between the variable calculated from the CEBQ "slowness in eating" scale and the eating speed variable estimated in minutes per day as well as between the 18-item and the 14-item questionnaires of adherence to the Mediterranean diet. All analyses were conducted using Stata 14 (StataCorp), and statistical significance was set at a 2-tailed P value <.05.

Results

Among the total of 1371 participants who attended the first visit, 49 were excluded for not meeting the inclusion criteria. Subsequently, 193 children had missing data for the time spent at breakfast, lunch, and/or dinner, precluding estimation of eating speed. In addition, 143 participants were excluded because of missing data or implausible reported energy intake. Finally, 30 participants with a current diagnosis of chronic diseases also were excluded from the analyses. Therefore, a total sample of 956 children (49% girls) were included in the analyses (**Figure**). For 88% of the study sample, the questionnaires were completed by the mother.

The main characteristics of the sample according to category of eating speed at baseline are shown in **Table III**. The mean age was 4.8 ± 1 years. Compared with participants in the slow-eating category, children with fast eating speed were more likely to have overweight or obesity, larger waist circumference, greater FMI, and higher systolic blood pressure and fasting plasma glucose concentration ($P < .05$). Mothers whose children were categorized as fast-eaters had higher BMI and weight status, as well as lower educational level ($P < .05$), compared with slow eaters. No other significant between-group differences were observed.




Figure. Flow diagram for the CORALS cohort.

Table IV lists the dietary characteristics according to eating speed categories. The percentage of total daily energy intake from carbohydrates differed significantly across the eating speed categories, higher in the fast-eating category. Participants in the fast-eating category also were more likely to have lower intakes of total fatty acids, monounsaturated fatty acids, and protein per body weight, as well as lower nut consumption and adherence to the Mediterranean diet.

The associations between eating speed category and outcome are shown in **Table V**. Compared with children in the slow-eating category, in unadjusted models and after adjusting for potential confounders, participants in the fast-eating category showed a higher risk for overweight/obesity (OR, 2.9; 95% CI, 1.8-4.4; $P < .01$). Compared with slower eaters, fast eaters had higher values for waist

circumference, FMI, systolic blood pressure, and fasting plasma glucose concentration, as well as lower diet quality (β , -0.5 ; 95% CI, -0.9 to -0.1 ; $P < .05$). When we excluded those children aged <5 years from the association with FMI, the results remained significant in the full adjusted model (β , 0.5 ; 95% CI, 0.2 - 0.8 ; $P < .01$). Compared with the slow-eating category, an inverse association was found between the fast-eating category and total cholesterol in the full adjusted model. No significant associations were observed between eating speed and the other cardiometabolic risk factors.

Data on eating speed measured by the CEBQ “slowness in eating” scale also showed positive associations between the fast-eating category and overweight/obesity (OR, 4.2; 95% CI, 2.7-6.4; $P < .01$), waist circumference (β , 3.1; 95% CI, 2.1-4.2; $P < .01$), and FMI (β , 0.7; 95% CI, 0.5-0.9; $P < .01$).

Table III. General characteristics of the study population according to eating speed categories

Characteristics	Eating speed categories			P value*
	Slow (>85 min), N = 291	Moderate (66–85 min), N = 325	Fast (<65 min), N = 340	
Eating speed, total min/day/3 main meals, mean \pm SD	108.3 \pm 18.3	76.2 \pm 4.9	53.8 \pm 9.2	<.001
Age, y, mean \pm SD	4.8 \pm 1.1	4.8 \pm 1.0	4.9 \pm 1.0	.100
Girls, % (n)	50.5 (147)	48.9 (159)	48.2 (164)	.845
Adiposity				
Weight status, % (n)				<.001
Underweight or normal weight	87.5 (251)	79.2 (251)	70.3 (234)	
Overweight or obesity	12.5 (36)	20.8 (66)	29.7 (99)	
Waist circumference, cm, mean \pm SD	50.3 \pm 7.5	52.4 \pm 6.0	53.4 \pm 8.3	<.001
FMI, kg/m ² , mean \pm SD	3.7 \pm 1.2	3.8 \pm 1.4	4.1 \pm 1.3	.003
Cardiovascular risk factors				
Systolic blood pressure, mmHg, mean \pm SD	102.0 \pm 12.9	103.1 \pm 13.2	105.2 \pm 12.9	.009
Diastolic blood pressure, mmHg, mean \pm SD	64.5 \pm 12.6	63.9 \pm 12.5	66.1 \pm 12.0	.068
Fasting glucose, mg/dL, mean \pm SD	75.8 \pm 8.8	77.8 \pm 8.7	78.8 \pm 9.9	<.001
Total cholesterol, mg/dL, median (IQR)	167 (149–186.5)	166 (149–184)	162 (145–179)	.055
HDL-c, mg/dL, median (IQR)	56 (48.5–66)	58 (50–67)	56.4 (49–64.7)	.495
LDL-c, mg/dL, median (IQR)	96 (83–112)	95 (83–112.6)	92 (78–105.1)	.083
Non-HDL-c, mg/dL, median (IQR)	107 (94–123.5)	107 (93–126)	104 (90–119)	.215
Triglycerides, mg/dL, median (IQR)	53 (44.5–67.5)	53 (42–65)	52 (43–66)	.556
Early life factors				
Weight at birth, kg, mean \pm SD	3.3 \pm 0.5	3.3 \pm 0.5	3.2 \pm 0.6	.554
Breastfeeding duration, % (n)				.872
<6 mo	40.3 (94)	42.7 (108)	41.6 (106)	
\geq 6 mo	59.7 (139)	57.3 (145)	58.4 (149)	
Mother's weight gain during pregnancy, kg, mean \pm SD	12.5 \pm 4.7	12.7 \pm 4.6	12.3 \pm 4.5	.589
Lifestyle factors				
Sports and physical activities, min/wk	197.2 \pm 117.7	179.0 \pm 105.1	178.7 \pm 115.2	.069
Total sleep duration per day, h	10.4 \pm 0.9	10.4 \pm 1.1	10.4 \pm 1.3	.667
Active lifestyle score, 0–13 points	9.2 \pm 1.6	9.1 \pm 1.6	9.1 \pm 1.6	.726
Maternal factors				
Age, y, mean \pm SD	40.7 \pm 4.5	41.4 \pm 7.4	40.6 \pm 5.8	.162
BMI, kg/m ² , mean \pm SD	24.3 \pm 4.6	25.1 \pm 4.8	25.1 \pm 5.4	.049
Weight status, % (n)				.022
Underweight or normal weight	67.7 (197)	56.9 (185)	61.2 (208)	
Overweight or obesity	32.3 (94)	43.1 (140)	38.8 (132)	
Educational level, % (n)				.021
Primary or lower	7.2 (21)	9.5 (31)	12.1 (41)	
Secondary	40.2 (117)	35.1 (114)	43.8 (149)	
Academic—graduated or underreported	52.6 (153)	55.4 (180)	44.1 (150)	
Socio-professional category, % (n)				.583
Homemaker/student/retired/unemployed	26.5 (77)	30.2 (98)	27.8 (93)	
Employee	73.5 (214)	69.9 (227)	72.2 (242)	

*P values calculated using the χ^2 test for categorical variables and ANOVA for continuous variables. A P value <.05 was considered significant (bold type).

No significant associations were observed for other cardiometabolic risk factors, but the direction of the associations was the same.

In correlation analyses, both eating speed variables (minutes/day and the CEBQ “slowness in eating” scale) were positively correlated ($r = 0.43$; $P < .001$), as well as the 2 questionnaires assessing adherence to the Mediterranean diet (the 18-item and 14-item questionnaires) ($r = 0.39$; $P < .01$).

Interaction analyses between eating speed category, sex, mother's BMI and educational level, weight at birth, and sedentary behaviors were not significant.

Discussion

Our results on eating speed and adiposity are in line with previous observational studies reporting that self-reported fast

eating speed is associated with higher weight status and larger waist circumference in adults^{28,29} and Asian children.^{12,30} In addition, a longitudinal study conducted in Japanese girls reported a positive association between eating speed and body fat mass percentage, but FMI was not assessed.³¹ Moreover, an 8-week interventional trial in American children that aimed to decelerate eating speed through nutritional education, timers, and interactive activities with families resulted in a decrease in eating speed in association with lower BMI.³²

The association between eating speed and blood pressure also has been explored in Asian adults, in which fast eating was associated with higher blood pressure.³³ Similarly in children, Yamagishi et al observed higher systolic blood pressure in boys with continuous fast-eating speed; the association was not significant in girls, however.³⁴ In line with our results regarding eating speed and fasting plasma glucose concentration, Nohara et al reported men, but not women, who

Table IV. Dietary characteristics according to eating speed category

Characteristics	Eating speed category			P value*
	Slow (>85 min), N = 291	Moderate (66-85 min), N = 325	Fast (<65 min), N = 340	
Dietary intake contribution				
Total energy intake, kcal/day, mean \pm SD	1698 \pm 340	1688 \pm 351	1714 \pm 370	.632
Carbohydrates, %, mean \pm SD	42.0 \pm 5.5	42.9 \pm 5.4	43.7 \pm 5.5	<.001
Proteins, %, mean \pm SD	15.1 \pm 2.1	15.0 \pm 2.2	14.9 \pm 2.0	.505
Total fatty acids, %, mean \pm SD	42.9 \pm 5.9	42.1 \pm 5.7	41.4 \pm 5.9	.007
Saturated fatty acids, %, mean \pm SD	14.1 \pm 1.9	14.0 \pm 2.0	13.8 \pm 2.1	.184
Monounsaturated fatty acids, %, mean \pm SD	19.6 \pm 4.7	18.9 \pm 4.4	18.4 \pm 4.4	.006
Polyunsaturated fatty acids, %, mean \pm SD	6.2 \pm 1.4	6.1 \pm 1.7	6.2 \pm 1.8	.941
Protein intake per body weight, g/kg/d, mean \pm SD	3.5 \pm 0.9	3.3 \pm 0.8	3.2 \pm 1.0	<.001
Fiber, g/1000 kcal, mean \pm SD	8.5 \pm 2.3	8.5 \pm 2.4	8.5 \pm 2.3	.985
Source goal, % (n)				.674
<14 g/1000 kcal	95.9 (279)	96.9 (315)	97.1 (330)	
\geq 14 g/1000 kcal	4.1 (12)	3.1 (10)	2.9 (10)	
Sodium, mg/dL, mean \pm SD	2308 \pm 736	2320 \pm 719	2307 \pm 684	.967
Diet quality				
Mediterranean Diet Score, 0-18 points	11.1 \pm 2.9	10.7 \pm 2.8	10.5 \pm 2.5	.015
Dairy products, mean \pm SD				
Milk, g/day	324.9 \pm 239.1	312.6 \pm 221.7	320.9 \pm 234.1	.795
Yogurt, g/day	105.0 \pm 79.2	111.2 \pm 85.7	117.6 \pm 110.9	.240
Cheese, g/day	12.5 \pm 10.8	12.3 \pm 12.9	12.6 \pm 12.3	.959
Derivative dairy products, g/day	80.2 \pm 82.9	85.5 \pm 93.9	90.4 \pm 99.5	.389
Protein foods, mean \pm SD				
White meat, g/day	24.7 \pm 9.8	23.9 \pm 9.0	23.9 \pm 9.2	.492
Unprocessed red meat, g/day	18.8 \pm 11.9	18.9 \pm 13.2	19.0 \pm 13.3	.974
Processed and derivatives meat products, g/day	26.3 \pm 16.6	25.7 \pm 14.4	24.5 \pm 14.7	.319
Egg, g/day	23.5 \pm 8.6	23.2 \pm 8.5	23.2 \pm 10.2	.889
Fish and seafood, g/day	34.9 \pm 17.0	33.9 \pm 17.4	32.5 \pm 17.9	.222
Vegetables and fruits, mean \pm SD				
Vegetables, g/day	76.8 \pm 52.8	77.4 \pm 51.1	75.8 \pm 52.7	.921
Tubers, g/day	40.3 \pm 18.6	39.9 \pm 19.2	41.4 \pm 20.3	.576
Fruits, g/day	188.6 \pm 113.2	190.0 \pm 124.1	181.4 \pm 122.9	.611
Nuts, g/day, mean \pm SD	4.2 \pm 5.7	3.3 \pm 4.4	3.5 \pm 4.6	.045
Cereals and legumes, mean \pm SD				
Legumes, g/day	14.0 \pm 6.4	14.0 \pm 6.2	14.3 \pm 8.5	.815
Refined bread, cereals, and pasta, g/day	70.3 \pm 32.2	72.7 \pm 34.9	74.5 \pm 38.4	.333
Brown bread, cereals, pasta and rice, g/day	5.5 \pm 13.5	3.7 \pm 10.7	4.5 \pm 11.2	.163
Miscellaneous, mean \pm SD				
Oil and fats, g/day	27.7 \pm 16.6	26.4 \pm 15.7	25.4 \pm 15.3	.212
Pastries, g/day	38.3 \pm 29.4	40.6 \pm 28.6	41.2 \pm 30.5	.454
Sugars and candies, g/day	12.9 \pm 10.5	13.2 \pm 10.6	13.6 \pm 11.0	.696
Beverages, mean \pm SD				
Water, g/day	852.6 \pm 390.9	859.0 \pm 371.8	348.9 \pm 363.2	.941
Sugary beverages, g/day	110.5 \pm 126.9	104.4 \pm 130.8	124.8 \pm 130.5	.111
Tea and infusions, g/day	5.3 \pm 28.3	6.0 \pm 24.7	8.8 \pm 32.9	.276

*P values calculated by the χ^2 test for categorical variables and ANOVA for continuous variables. P values <.05 were considered significant (bold type).

reported fast-eating speed had higher fasting glucose concentrations and blood pressure values.³⁵ Another study of middle-aged Japanese adults reported an independent and positive association between fast eating and insulin resistance; however, the association with blood glucose level was not assessed.³⁶ To date, no published studies have explored the association between eating speed and fasting plasma glucose in children. In contrast with a recent review that reported significant associations between eating speed and triglyceride and lower HDL-c levels in adults,² our results did not show any significant associations in a sample of children. Inconclusive results in these studies may be related to variations in populations (eg, ethnicity, age, culture), as well as additional differences in methodology.

In the present study, we observed an inverse association between eating speed and the adherence to Mediterranean

diet. This result could be explained by the main characteristics of the Mediterranean diet, such as its high content of plant-based foods (which provide high amounts of dietary fiber) and fish/seafood and low content of red meat/processed meat products.³⁷ Even though there were no significant differences in the majority of food groups in our analyses, the overall beneficial effect of the Mediterranean diet is greater than the effects of its individual components. Furthermore, in a meta-analysis³⁸ and a randomized-crossover study,³⁹ positive effects were observed between the consumption of fiber or protein and appetite suppression as well as satiety, which might have an impact on eating speed.⁷ Additionally, associations between faster eating,¹⁶ sweet-tasting foods,¹⁰ and ultra-processed foods⁴⁰ have been reported. Ultra-processed foods are characterized by their high energy, salt, sugar, and fat content,⁴¹ which

Table V. Crude and multivariate β -coefficients (95% CI) or OR (95% CI) of adiposity, diet quality, and cardiometabolic risk factors according to eating speed categories in the total population

Variables	Eating speed category		
	Slow (>85 min)	Moderate (66-85 min)	Fast (<65 min)
Adiposity			
Weight status, n	287	317	333
Crude model	1 (ref)	1.8 (1.2-2.9) [†]	2.9 (1.9-4.5) [†]
Model 1	1 (ref)	1.8 (1.1-2.8)*	2.8 (1.8-4.4) [†]
Model 2	1 (ref)	1.8 (1.1-2.8)*	2.9 (1.8-4.4) [†]
Waist circumference, n	285	323	338
Crude model	0 (ref)	1.9 (0.7-3.0) [†]	2.9 (1.8-4.0) [†]
Model 1	0 (ref)	1.7 (0.6-2.7) [†]	2.5 (1.4-3.6) [†]
Model 2	0 (ref)	1.7 (0.6-2.8) [†]	2.5 (1.4-3.8) [†]
FMI, n	258	283	303
Crude model	0 (ref)	0.2 (−0.1 to 0.4)	0.4 (0.2-0.6) [†]
Model 1	0 (ref)	0.1 (−0.1 to 0.3)	0.3 (0.1-0.5) [†]
Model 2	0 (ref)	0.1 (−0.1 to 0.3)	0.3 (0.1-0.5) [†]
Cardiovascular risk factors			
Systolic blood pressure, n	274	304	325
Crude model	0 (ref)	1.0 (−1.1 to 3.2)	3.2 (1.1-5.3) [†]
Model 1	0 (ref)	0.8 (−1.3 to 2.9)	2.8 (0.7-4.9) [†]
Model 2	0 (ref)	0.8 (−1.3 to 3.0)	2.8 (0.6-4.9)*
Diastolic blood pressure, n	275	305	325
Crude model	0 (ref)	−0.6 (−2.6 to 1.4)	1.6 (−0.4 to 3.6)
Model 1	0 (ref)	−0.7 (−2.7 to 1.3)	1.6 (−0.4 to 3.6)
Model 2	0 (ref)	−0.8 (−2.8 to 1.2)	1.5 (−0.5 to 3.5)
Fasting glucose, n	241	270	290
Crude model	0 (ref)	2.0 (0.4-3.6)*	3.0 (1.5-4.6) [†]
Model 1	0 (ref)	2.2 (0.7-3.7) [†]	2.8 (1.3-4.3) [†]
Model 2	0 (ref)	2.1 (0.6-3.6) [†]	2.7 (1.2-4.2) [†]
Total cholesterol, n	240	271	287
Crude model	0 (ref)	−0.5 (−5.4 to 4.4)	−4.5 (−9.4 to 0.3)
Model 1	0 (ref)	−0.2 (−5.1 to 4.6)	−4.5 (−9.4 to 0.3)
Model 2	0 (ref)	−0.6 (−5.4 to 4.3)	−5.0 (−9.8 to −0.2)*
HDL-c, n	240	270	286
Crude model	0 (ref)	0.0 (−2.3 to 2.4)	−1.0 (−3.3 to 1.3)
Model 1	0 (ref)	0.2 (−2.2 to 2.5)	−0.9 (−3.3 to 1.4)
Model 2	0 (ref)	0.2 (−2.1 to 2.6)	−0.9 (−3.2 to 1.5)
LDL-c, n	225	255	273
Crude model	0 (ref)	0.3 (−4.1 to 4.7)	−3.2 (−7.4 to 1.0)
Model 1	0 (ref)	0.3 (−4.1 to 4.6)	−3.2 (−7.5 to 1.0)
Model 2	0 (ref)	0.1 (−4.2 to 4.5)	−3.7 (−7.9 to 0.6)
Non-HDL-c, n	240	270	286
Crude model	0 (ref)	−1.0 (−5.4 to 3.4)	−3.5 (−7.8 to 0.9)
Model 1	0 (ref)	−0.8 (−5.2 to 3.6)	−3.5 (−7.9 to 0.9)
Model 2	0 (ref)	−1.2 (−5.6 to 3.2)	−4.1 (−8.4 to 0.3)
Triglycerides, n	240	271	286
Crude model	0 (ref)	−1.6 (−5.3 to 2.1)	−2.3 (−6.0 to 1.3)
Model 1	0 (ref)	−2.3 (−5.9 to 1.2)	−3.4 (−7.0 to 0.1)
Model 2	0 (ref)	−2.4 (−6.0 to 1.2)	−3.5 (−7.1 to 0.1)
Diet quality			
Mediterranean Diet Score, points	289	324	339
Crude model	0 (ref)	−0.3 (−0.8 to 0.1)	−0.6 (−1.1 to −0.2) [†]
Model 1	0 (ref)	−0.3 (−0.7 to 0.1)	−0.5 (−0.9 to −0.1)*
Model 2	0 (ref)	−0.3 (−0.7 to 0.1)	−0.5 (−0.9 to −0.1)*

Multiple linear regressions models were fitted and expressed in β coefficients (95% CI). For weight status, logistic regressions models were fitted and expressed in OR (95% CI). The weight status is a dichotomous outcome (underweight or normal weight -1- and overweight or obesity -2-).

Model 1 adjusted by center, sex, age, maternal BMI, and mother's educational level.

Model 2 adjusted additionally by total energy intake, breastfeeding, total minutes of physical activity per week, and total hours of sleep per day.

*Associations with $P < .05$.

†Associations with $P < .01$.

may contribute to food reinforcement and a higher overall energy intake,⁴¹ perhaps because of their high palatability.⁴⁰ On the other hand, an unexpected result in our analyses was the association between fast eating speed and lower total cholesterol. In this sense, a potential association between

lower fat intake and a decrease in total cholesterol and LDL-c has been reported,⁴² which could partially explain this result, as our analyses showed a lower percentage of total fatty acids in those participants in the fast-eating category.

This study has some limitations that merit mention. First, because it is a cross-sectional study, we cannot draw cause-and-effect conclusions or discard bidirectional associations. Second, our study sample included Spanish preschool children, and thus our findings cannot be extrapolated to other populations. Third, we cannot dismiss the possibility that associations might be related to residual confounding or that undetected cardiometabolic disorders in our study population may exist because of age. Fourth, equations for bioelectrical impedance analysis have not been designed for children aged <5 years; however results from the analyses excluding children aged 3-4 years remained similar to those in the total population. Fifth, although eating speed was self-reported, it was estimated according to the time reported by main caregivers, not based on self-perception. Moreover, analyses from the CEBQ “slowness in eating” scale showed similar results as those observed in the main analyses for adiposity. Sixth, the 18-item Mediterranean diet questionnaire has not been validated in children; nevertheless, it was positively correlated to the validated 14-item Mediterranean diet questionnaire, and results should be interpreted cautiously.

In conclusion, eating fast is associated with higher levels of adiposity and certain cardiometabolic risk factors, as well as lower adherence to the Mediterranean diet in Spanish children. However, further long-term studies and clinical trials are needed to confirm these associations. ■

We thank all the CORALS participants and their parents or caregivers as well as the health centers and primary schools for their collaboration, the CORALS personnel for their outstanding support, and the staff of all associated primary care centers for their exceptional work. We particularly acknowledge the Institut d'Investigacion Sanitaria Pere Virgili (IISPV) Biobank (PT20/00197), which is integrated in the Instituto de Salud Carlos III (ISCIII) Platform for Biobanks and Biomodel.

Submitted for publication Apr 19, 2022; last revision received Aug 15, 2022; accepted Aug 18, 2022.

Reprint requests: Nancy Babio, PhD, Unitat de Nutrició Humana, Departament de Bioquímica i Biotecnologia, Universitat Rovira i Virgili, Reus, Spain. E-mail: nancy.babio@urv.cat

Data Statement

Data sharing statement available at www.jpeds.com.

References

- di Cesare M, Sorić M, Bovet P, Miranda JJ, Bhutta Z, Stevens GA, et al. The epidemiological burden of obesity in childhood: a worldwide epidemic requiring urgent action. *BMC Med* 2019;17:212. <https://doi.org/10.1186/s12916-019-1449-8>
- Garcidueñas-Fimbres TE, Paz-Graniel I, Nishi SK, Salas-Salvadó J, Babio N. Eating speed, eating frequency, and their relationships with diet quality, adiposity, and metabolic syndrome, or its components. *Nutrients* 2021;13:1687. <https://doi.org/10.3390/nu13051687>
- Okubo H, Murakami K, Masayasu S, Sasaki S. The relationship of eating rate and degree of chewing to body weight status among preschool children in Japan: a nationwide cross-sectional study. *Nutrients* 2018;11:64. <https://doi.org/10.3390/nu11010064>
- Kolay E, Bykowska-Derda A, Abdulsamad S, Kaluzna M, Samarzewska K, Ruchala M, et al. Self-reported eating speed is associated with indicators of obesity in adults: a systematic review and meta-analysis. *Healthcare (Basel)* 2021;9:1559. <https://doi.org/10.3390/healthcare9111559>
- Paz-Graniel I, Babio N, Mendez I, Salas-Salvadó J. Association between eating speed and classical cardiovascular risk factors: a cross-sectional study. *Nutrients* 2019;11:83. <https://doi.org/10.3390/nu11010083>
- Hamada Y, Miyaji A, Hayashi Y, Matsumoto N, Nishiwaki M, Hayashi N. Objective and subjective eating speeds are related to body composition and shape in female college students. *J Nutr Sci Vitaminol (Tokyo)* 2017;63:174-9. <https://doi.org/10.3177/jnsv.63.174>
- Kokkinos A, le Roux CW, Alexiadou K, Tentolouris N, Vincent RP, Kyriaki D, et al. Eating slowly increases the postprandial response of the anorexigenic gut hormones, peptide YY and glucagon-like peptide-1. *J Clin Endocrinol Metab* 2010;95:333-7. <https://doi.org/10.1210/jc.2009-1018>
- Ekuni D, Furuta M, Takeuchi N, Tomofuji T, Morita M. Self-reports of eating quickly are related to a decreased number of chews until first swallow, total number of chews, and total duration of chewing in young people. *Arch Oral Biol* 2012;57:981-6. <https://doi.org/10.1016/j.archoralbio.2012.02.001>
- Spiegel TA, Wadden TA, Foster GD. Objective measurement of eating rate during behavioral treatment of obesity. *Behav Ther* 1991;22:61-7. [https://doi.org/10.1016/S0005-7894\(05\)80244-8](https://doi.org/10.1016/S0005-7894(05)80244-8)
- van den Boer J, Werts M, Siebelink E, de Graaf C, Mars M. The availability of slow and fast calories in the Dutch diet: the current situation and opportunities for interventions. *Foods* 2017;6:87. <https://doi.org/10.3390/foods6100087>
- Forde CG, van Kuijk N, Thaler T, de Graaf C, Martin N. Oral processing characteristics of solid savoury meal components, and relationship with food composition, sensory attributes and expected satiation. *Appetite* 2013;60:208-19. <https://doi.org/10.1016/j.appet.2012.09.015>
- Fogel A, Goh AT, Fries LR, Sadananthan SA, Velan SS, Michael N, et al. Faster eating rates are associated with higher energy intakes during an ad libitum meal, higher BMI and greater adiposity among 4-5-year-old children: results from the Growing Up in Singapore Towards Healthy Outcomes (GUSTO) cohort. *Br J Nutr* 2017;117:1042-51. <https://doi.org/10.1017/S0007114517000848>
- Viskaal-van Dongen M, Kok FJ, de Graaf C. Eating rate of commonly consumed foods promotes food and energy intake. *Appetite* 2011;56:25-31. <https://doi.org/10.1016/j.appet.2010.11.141>
- Forde CG, Leong C, Chia-Ming E, McCrickerd K. Fast or slow-foods? Describing natural variations in oral processing characteristics across a wide range of Asian foods. *Food Funct* 2017;8:595-606. <https://doi.org/10.1039/c6fo01286h>
- Poti JM, Mendez MA, Ng SW, Popkin BM. Is the degree of food processing and convenience linked with the nutritional quality of foods purchased by US households? *Am J Clin Nutr* 2015;101:1251-62. <https://doi.org/10.3945/ajcn.114.100925>
- Teo PS, van Dam RM, Forde CG. Combined impact of a faster self-reported eating rate and higher dietary energy intake rate on energy intake and adiposity. *Nutrients* 2020;12:3264. <https://doi.org/10.3390/nu12113264>
- Carnell S, Wardle J. Measuring behavioural susceptibility to obesity: validation of the child eating behaviour questionnaire. *Appetite* 2007;48:104-13. <https://doi.org/10.1016/j.appet.2006.07.075>
- Jimeno-Martínez A, Maneschy I, Moreno LA, Bueno-Lozano G, de Miguel-Etayo P, Flores-Rojas K, et al. Reliability and validation of the Child Eating Behavior Questionnaire in 3- to 6-year-old Spanish children. *Front Psychol* 2022;13:705912. <https://doi.org/10.3389/fpsyg.2022.705912>
- Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 2012;7:284-94. <https://doi.org/10.1111/j.2047-6310.2012.00064.x>
- VanItallie TB, Yang MU, Heymsfield SB, Funk RC, Boileau RA. Height-normalized indices of the body's fat-free mass and fat mass: potentially useful indicators of nutritional status. *Am J Clin Nutr* 1990;52:953-9. <https://doi.org/10.1093/ajcn/52.6.953>

21. Frost PH, Havel RJ. Rationale for use of non-high-density lipoprotein cholesterol rather than low-density lipoprotein cholesterol as a tool for lipoprotein cholesterol screening and assessment of risk and therapy. *Am J Cardiol* 1998;81:26B-31B. [https://doi.org/10.1016/s0002-9149\(98\)00034-4](https://doi.org/10.1016/s0002-9149(98)00034-4)
22. Schröder H, Fitó M, Estruch R, Martínez-González MA, Corella D, Salas-Salvado J, et al. A short screener is valid for assessing Mediterranean diet adherence among older Spanish men and women. *J Nutr* 2011;141:1140-5. <https://doi.org/10.3945/jn.110.135566>
23. Verbestel V, De Henauw S, Bammann K, Barba G, Hadjigeorgiou C, Eiben G, et al. Are context-specific measures of parental-reported physical activity and sedentary behaviour associated with accelerometer data in 2-9-year-old European children? *Public Health Nutr* 2015;18:860-8. <https://doi.org/10.1017/S136898001400086X>
24. Fernández-Ballart JD, Piñol JL, Zazpe I, Corella D, Carrasco P, Toledo E, et al. Relative validity of a semi-quantitative food-frequency questionnaire in an elderly Mediterranean population of Spain. *Br J Nutr* 2010;103:1808-16. <https://doi.org/10.1017/S0007114509993837>
25. Farran A, Zamora R, Cervera P, Centre d'Enseymament Superior de Nutrició i Dietètica. *Tablas de composición de alimentos del CESNID*. 3rd ed. Universitat de Barcelona: McGraw-Hill Interamericana; 2004 [in Spanish].
26. Börnhorst C, Huybrechts I, Ahrens W, Eiben G, Michels N, Pala V, et al. Prevalence and determinants of misreporting among European children in proxy-reported 24 h dietary recalls. *Br J Nutr* 2013;109:1257-65. <https://doi.org/10.1017/S0007114512003194>
27. de Waal T, Pannekoek J, Scholtus S. *Statistical data editing and imputation*. Hoboken (NJ): John Wiley & Sons; 2011.
28. Yamane M, Ekuni D, Mizutani S, Kataoka K, Sakumoto-Kataoka M, Kawabata Y, et al. Relationships between eating quickly and weight gain in Japanese university students: a longitudinal study. *Obesity* (Silver Spring) 2014;22:2262-6. <https://doi.org/10.1002/oby.20842>
29. Zhu B, Haruyama Y, Muto T, Yamazaki T. Association between eating speed and metabolic syndrome in a three-year population-based cohort study. *J Epidemiol* 2015;25:332-6. <https://doi.org/10.2188/jea.JE20140131>
30. Okubo H, Miyake Y, Sasaki S, Tanaka K, Hirota Y. Rate of eating in early life is positively associated with current and later body mass index among young Japanese children: the Osaka Maternal and Child Health Study. *Nutr Res* 2017;37:20-8. <https://doi.org/10.1016/j.nutres.2016.11.011>
31. Ochiai H, Shirasawa T, Ohtsu T, Nishimura R, Morimoto A, Hoshino H, et al. The impact of eating quickly on anthropometric variables among schoolgirls: a prospective cohort study in Japan. *Eur J Public Health* 2014;24:691-5. <https://doi.org/10.1093/eurpub/ckt120>
32. Faith MS, Diewald LK, Crabbe S, Burgess B, Berkowitz RI. Reduced eating pace (RePace) behavioral intervention for children prone to or with obesity: does the turtle win the race? *Obesity* (Silver Spring) 2019;27:121-9. <https://doi.org/10.1002/oby.22329>
33. Tao L, Yang K, Huang F, Liu X, Li X, Luo Y, et al. Association between self-reported eating speed and metabolic syndrome in a Beijing adult population: a cross-sectional study. *BMC Public Health* 2018;18:855. <https://doi.org/10.1186/s12889-018-5784-z>
34. Yamagishi K, Sairenchi T, Sawada N, Sunou K, Sata M, Murai U, et al. Impact of speed-eating habit on subsequent body mass index and blood pressure among schoolchildren the Ibaraki Children's Cohort Study (IBA-CHIL). *Circ J* 2018;82:419-22. <https://doi.org/10.1253/circj.CJ-17-0287>
35. Nohara A, Maejima Y, Shimomura K, Kumamoto K, Takahashi M, Akuzawa M, et al. Self-awareness of fast eating and its impact on diagnostic components of metabolic syndrome among middle-aged Japanese males and females. *Endocr Regul* 2015;49:91-6. https://doi.org/10.4149/endo_2015_02_91
36. Otsuka R, Tamakoshi K, Yatsuya H, Wada K, Matsushita K, OuYang P, et al. Eating fast leads to insulin resistance: findings in middle-aged Japanese men and women. *Prev Med* 2008;46:154-9. <https://doi.org/10.1016/j.ypmed.2007.07.031>
37. Davis C, Bryan J, Hodgson J, Murphy K. Definition of the Mediterranean Diet; a literature review. *Nutrients* 2015;7:9139-53. <https://doi.org/10.3390/nu7115459>
38. Qiu M, Zhang Y, Long Z, He Y. Effect of protein-rich breakfast on subsequent energy intake and subjective appetite in children and adolescents: systematic review and meta-analysis of randomized controlled trials. *Nutrients* 2021;13:2840. <https://doi.org/10.3390/nu13082840>
39. Halliday TM, Liu SV, Moore LB, Hedrick VE, Davy BM. Adolescents perceive a low added sugar adequate fiber diet to be more satiating and equally palatable compared to a high added sugar low fiber diet in a randomized-crossover design controlled feeding pilot trial. *Eat Behav* 2018;30:9-15. <https://doi.org/10.1016/j.eatbeh.2018.05.004>
40. Forde CG, Mars M, de Graaf K. Ultra-processing or oral processing? A role for energy density and eating rate in moderating energy intake from processed foods. *Curr Dev Nutr* 2020;4:nzaa019. <https://doi.org/10.1093/cdn/nzaa019>
41. Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, et al. Ultra-processed diets cause excess calorie intake and weight gain: an inpatient randomized controlled trial of ad libitum food intake. *Cell Metab* 2019;30:67-77.e3. <https://doi.org/10.1016/j.cmet.2019.05.008>
42. Naude CE, Visser ME, Nguyen KA, Durao S, Schoonees A. Effects of total fat intake on bodyweight in children. *Cochrane Database Syst Rev* 2018;7:CD012960. <https://doi.org/10.1002/14651858.CD012960.pub2>

Table I. The 18-item questionnaire of adherence to the Mediterranean diet in children

Questions	Criterion for 1 point
1. Do you use extra-virgin olive oil as the main culinary fat?	Yes
2. Do you consume 3 or more tablespoons of olive oil per day (including that used for frying, salads or in meals away from home)? (1 tbsp=10 ml)	Yes
3. Do you consume 2 or more servings of vegetables per day? During the week, do you eat any of these servings as raw vegetables or salad? (1 serving= 50-80 g)	Yes
4. Do you consume 3 or more small fruits per day? (small fruit = 100 g)	Yes
5. Do you consume whole grains (bread, cereals, pasta, or rice) 3 or more times a week, instead of refined grains?	Yes
6. Do you consume at least 1 serving per day of fermented milk, plain yogurt or goat's or sheep's cheese? (1 commercial portion of fermented milk or yogurt or 25 g of cheese)	Yes
7. Do you consume 2-3 or more servings a week of legumes? (1 serving = 40 g raw weight)	Yes
8. Do you consume 3 or more servings a week of fish/seafood? (1 serving = 40-70 g)	Yes
9. Do you consume at least 3 servings a week of nuts? (1 serving = 15-20 g)	Yes
10. Do you consume preferably chicken, turkey, or rabbit instead of beef, pork, hamburgers, or sausage?	Yes
11. Do you consume 2 or more times a week cooked vegetables, pasta, rice, or other dishes seasoned with tomato, garlic, onion, or leek sauce simmered with olive oil (sofrito)?	Yes
12. Do you eat red meat, hamburgers, sausages, or derivative/processed meat products less than 2 times a week?	Yes
13. Do you eat less than 1 serving per day of butter or cream? (1 serving = 12 g)	Yes
14. Do you drink less than 2 glasses a week of carbonated and/or sugary beverages (sodas, coke, juices, nectars)?	Yes
15. Do you consume chips, gummy worms, sweets less than 1 time a week?	Yes
16. Do you consume dairy desserts, such as custard, ice cream, dairy smoothies, petit Suisse, vegetable drinks, etc, less than 1 time a week?	Yes
17. Do you consume pastries, stuffed cookies, sweets, or cakes less than 2 times a week?	Yes
18. Do you consume precooked or ready-to-eat foods less than 1 time a week?	Yes

Table II. Questionnaire for evaluating an active lifestyle in children

Questions	Criterion for 1 point
19. Does she/he walk or cycle to the school at least 3 days a week?	Yes
20. Does she/he play outdoors such as in parks or playgrounds, during leisure time, at least 3 days a week?	Yes
21. Does she/he participate in extracurricular sports or recreational activities like basketball, soccer, athletics, cycling, swimming, rhythmic gymnastics, other sports, traditional games, etc, at least 2 days a week?	Yes
22. Does she/he do strength and flexibility activities like dancing, yoga, or martial arts at least 4 days a week?	Yes
23. Does she/he preferentially use the stairs, such as at home?	Yes
24. Does she/he walk every day for at least 30 minutes, even if it is not continuous?	Yes
25. Does she/he participate in outdoors activities with the family during weekdays at least 1 time per week? (cycling, playing ball)	Yes
26. Does she/he participate in outdoors activities like family excursions during the weekends, at least 2 times per month?	Yes
27. During the leisure time, does she/he participate in activities that require more movement such as dancing, running, jumping, instead of activities like painting, sitting down to play, using the computer, etc?	Yes
28. Does she/he like the physical education classes at school?	Yes
29. During breaks at school, does she/he prefer activities like running, jumping or other activities instead to nonmoving games (talking, sitting down, playing with cars or dolls, etc)?	Yes
30. During weekdays, does she/he use audiovisual media, such as television, computers, tablets, or cell phones, less than 2 hours a day?	Yes
31. During weekends, does she/he use audiovisual media, such as television, computers, tablets, or cell phones, less than 2 hours a day?	Yes